arXiv:gr-qc/0105119v1 30 May 2001

The Kretschmann scalar for 5D Vacua

Kayll Lake[1]

Department of Physics, Queen's University
Kingston, Ontario, K7L 3N6, Canada

We present a streamlined calculation of the Kretschmann scalar for 5D Vacua considered recently by Fukui *et al* .

I. INTRODUCTION

Recently, Fukui *et al* [1] examined the 5D metric

$$ds^2 = \sigma\{dt^2 - a^2[\frac{dr^2}{(1-kr^2)} + r^2 d\Omega^2]\} + \epsilon b^2 dl^2, \tag{1}$$

where $k = \pm 1, 0$, $d\Omega^2 \equiv d\theta^2 + \sin^2\theta\, d\phi^2$, $\epsilon = \pm 1$, $\sigma = \sigma(l)$, $a = a(t,l)$ and $b = b(t,l)$, with regard to the 5D vacuum condition $R_{AB} = 0$. The authors comment that the associated 5D Kretschmann curvature invariant $K \equiv R^{ABCD}R_{ABCD}$ is difficult to calculate even by computer. The purpose of this note is to show how to trivialize this calculation.

II. NOTATION AND KRETSCHMANN DECOMPOSITION

Write R_{ABCD} for the Riemann tenor, R_{AB} for the Ricci tensor, R for the Ricci scalar ($\equiv R^A_A$) and C_{ABCD} for the Weyl tensor where the dimension of the space is n. A straightforward calculation gives the identity

$$R_{ABCD}R^{ABCD} = \frac{8}{(n-2)^2}R_{AB}R^{AB} + \frac{4(n^2-8n+13)}{(n-1)^2(n-2)^2}R^2 + C_{ABCD}C^{ABCD} \tag{2}$$

so that for vacua we need consider only the Weyl invariant $C_{ABCD}C^{ABCD}$ which, for the metric (1), is given by

$$\frac{(2\,aa_{ll}b^3\epsilon - 2\,a_l^2b^3\epsilon - 2\,kb^3\epsilon + 2\,aa_tb_tb^2\epsilon + a\sigma_l ba_l - 2\,ab_l\sigma a_l + 2\,b\sigma aa_{ll} - 2\,\epsilon\, b^2b_{tt}a^2 - 2\,b\sigma a_l^2)^2}{2\sigma^2b^6a^4} \tag{3}$$

where a subscript denotes differentiation. The calculation proceeds in about $\frac{1}{5}$ second on a contemporary PC under GRTensorII [2]. It turns out that for metrics of the form (1), prior to the vacuum requirement, $R_{AB}R^{AB}$ is cumbersome. Even the Ricci scalar turns out to be larger than the Weyl invariant calculated here. Under the restrictions (11) and (12) in Fukui *et al* [1] (3) reduces to their equation (13).

ACKNOWLEDGMENTS

This work was supported by a grant from the Natural Sciences and Engineering Research Council of Canada.

---

[1] E-mail: `lake@astro.queensu.ca`

[1] T. Fukui, S. Seahra and P. S. Wesson, gr-qc/0105112.

[2] Portions of this work were made possible by use of *GRTensorII*. This is a package which runs within Maple. It is entirely distinct from packages distributed with Maple and must be obtained independently. The GRTensorII software and documentation is distributed freely on the World-Wide-Web from the address `http://grtensor.phy.queensu.ca`. Further details concerning the calculation reported here can be found at `http://grtensor.phy.queensu.ca/five5`.